

**Supplemental Table – Baseline and Procedural Characteristics of all Patients, Sorted by Stent Type**

Variable	Paclitaxel	Sirolimus	Everolimus	P
	eluting stent (n=1298)	eluting stent (n=2599)	eluting stent (n=1466)	
<b>Age (years)</b>	63.9±11.6	62.7±11.5	64.9±11.7	<0.0001
<b>Male sex</b>	990 (76.3)	1976 (76.0)	1118 (76.3)	0.98
<b>BMI</b>	27.0±4.1	27.2±4.3	27.2±4.4	0.12
<b>Hypertension</b>	748 (57.6)	1422 (54.7)	857 (59.3)	0.01
<b>Family history of CAD</b>	334 (26.5)	738 (28.4)	420 (29)	0.30
<b>Current smoking</b>	671 (51.7)	1371 (52.8)	736 (50.9)	0.51
<b>Dyslipidemia</b>	683 (52.6)	1403 (54.0)	794 (54.9)	0.48
<b>Diabetes Mellitus</b>	210 (16.2)	471 (18.1)	246 (17.0)	0.30
<b>Renal failure (GFR&lt;60mL/min)</b>	138 (14.2)	294 (15.6)	146 (11.6)	0.005
<b>Acute coronary syndrome</b>	714 (55.1)	1385 (53.3)	855 (58.3)	0.009
Unstable angina	67 (9.4)	129 (9.3)	108 (12.6)	
Non-ST-segment-elevation MI	292 (40.9)	622 (44.9)	382 (44.7)	
ST-segment-elevation MI	355 (49.7)	634 (45.8)	365 (42.7)	
<b>Cardiogenic shock</b>	9 (0.7)	28 (1.1)	33 (2.3)	0.001
<b>LVEF</b>	55.3±11.5	54.3±11.9	51.5±12.7	<0.0001
Moderate-severely impaired LVEF	194 (14.9)	449 (17.3)	370 (25.2)	<0.0001
Mildly impaired LVEF	300 (23.0)	584 (22.4)	360 (24.6)	
Normal LVEF	804 (61.9)	1566 (60.3)	736 (50.2)	
<b>Multivessel treatment</b>	218 (16.8)	407 (15.7)	342 (23.3)	<0.0001
<b>Lesions treated per patient</b>	1.5±0.7	1.5±0.7	1.8±1.0	<0.0001
<b>Culprit left main coronary artery</b>	30 (2.3)	46 (1.8)	67 (4.6)	<0.0001
<b>Arterial bypass graft</b>	5 (0.4)	5 (0.2)	3 (0.2)	0.49
<b>Saphenous vein graft</b>	33 (2.5)	77 (3.0)	47 (3.2)	0.58
<b>Multistent treatment</b>	614 (47.3)	1215 (46.7)	844 (57.6)	<0.0001
<b>Average stent diameter</b>	2.9±0.4	2.9±0.6	2.9±0.4	0.21
<b>Total stent length per patient</b>	29.1±16.9	31.9±18.3	32.8±18.9	<0.0001
<b>Glycoprotein IIb/IIIa antagonist</b>	245 (18.9)	501 (19.3)	438 (29.9)	<0.0001
<b>Aspirin at discharge</b>	1282 (98.8)	2549 (98.1)	1433 (97.7)	0.13
<b>Clopidogrel at discharge</b>	1290 (99.4)	2566 (98.7)	1432 (97.7)	0.001
<b>Oral anticoagulation at discharge</b>	15 (1.2)	51 (2.0)	27 (1.8)	0.17

Categorical variables are expressed as count (percentage). Valid percentages may vary for some counts, because of missing values. Continuous variables are expressed as mean ±SD when appropriate. BMI indicates body mass index; CAD, coronary artery disease; GFR, glomerular filtration rate; LVEF, left ventricular ejection fraction; MI, myocardial infarction.