

Supplemental table 4: Patient kidney characteristics					
	All (213)	Men (83)		Women (130)	
		Classical (57)	Non-classical (26)	Classical (98)	Non-classical (32)
KDIGO CKD stages**					
Patients with a known CKD stage, n (%)	152 (71%)	41 (72%)	13 (50%)	82 (84%)	16 (50%)
G1 (eGFR >90), n (%)	75 (49%)	23 (56%)	3 (23%)	37 (45%)	12 (75%)
G2 (eGFR 60-89), n (%)	48 (32%)	9 (22%)	4 (31%)	33 (40%)	2 (13%)
G3a (eGFR 45-59), n (%)	13 (9%)	1 (2%)	2 (15%)	8 (10%)	2 (13%)
G3b (eGFR 30-44), n (%)	12 (8%)	7 (17%)	2 (15%)	3 (4%)	0 (0%)
G4 (eGFR 15-29), n (%)	3 (2%)	1 (2%)	1 (8%)	1 (1%)	0 (0%)
G5 (eGFR <15), n (%)	1 (1%)	0 (0%)	1 (8%)	0 (0%)	0 (0%)
Proteinuria (microalbuminuria > 300 milligrams/ 24 hours)**					

Patients with a known proteinuria status, n (%)	71 (33%)	16 (28%)	7 (27%)	41 (42%)	7 (22%)
Present proteinuria, n (%)	12 (17%)	4 (25%)	3 (43%)	5 (12%)	0 (0%)
Absent proteinuria, n (%)	59 (83%)	12 (75%)	4 (57%)	36 (88%)	7 (100%)
<p>KDIGO= Kidney Disease Improving Global Outcomes; CKD= Chronic Kidney Disease; eGFR= estimated Glomerular Filtration Rate, according to the Chronic Kidney Disease Epidemiology Collaboration equation and measured in milliliters/ minute per 1.73 meter²</p> <p>** The KDIGO CKD and proteinuria status are measurements, which are known at the time of the last follow-up (with a minimum and maximum range of 1 year between the measurement and the last follow-up date).</p>					